

Rheumatology Enrollment Form (R-Z)

www.pyramidspharmacy.com

All our referral forms are available on our website.

10970 Shadow Creek Pkwv. Suite 110.1 I Pearland. TX 77584 I Phone: 1.888.375.1920 I Fax: 1.888.383.2140

1: Patient Information					
Patient Name:		Birthdate:	Sex: Male Female Height:	Weight: □lbs □kg	
Soc. Sec. #: Prefe			· ·	· · ·	
Address:			City:		
Insurance Information: Please fax FRONT and BACK copy of ALL Insurance cards (Prescription and Medical)					
Plan Name:Plan Phone:					
2: Prescriber Information					
Prescriber Name:			DEA#: NPI#:	Tax ID#:	
Address:			Phone: ()	Fax: (
City:		State: Zip:	Key Contact:	Phone: (
3: Diagnosis/Clinical Information Please FAX recent clinical notes, Labs, Tests, with the prescription to expedite the Prior Authorization					
Diagnosis (ICD-10): ☐ M06.9 Rheumatoid Arthritis, Unspecified ☐ M45.9 Ankylosing Spondylitis ☐ M08.00 Juvenile Rheumatoid Arthritis ☐ L40.50 Arthropathic Psoriasis ☐ Other:		TB Test Completed: ☐ Yes ☐ No Date of Negative Test: If history of latent TB, has patient retreatment? ☐ Yes ☐ No Hep B Screening: ☐ Positive ☐ Negative ☐ N/A Date of Hep B Test:	Please list previously tried and failed therapies & reason for discontinuing: Please list previously tried and failed therapies & reason for discontinuing: Azathioprine: Corticosteroids: Leflunomide: Methotrexate: NSAIDs: Hydroxychloroquine:		
4: Prescription Information					
Medication	Strength		Directions	Quantity Refills	
□ Remicade®	□ 100 mg Vial Patient Dosing Wt: _kg	maintenance dose. Maintenance Dose: Infu	mg/kg IV at Weeks 0, 2, and 6, then use mg/kg IV every 6 weeks. use mg/kg IV every 8 weeks.		
☐ Renflexis®	□ 100 mg Vial Patient Dosing Wt:kg	□ Initial Dose: Infuse maintenance dose. □ Maintenance Dose: Infu	mg/kg IV at Weeks 0, 2, and 6, then lse mg/kg IV every 6 weeks. lse mg/kg IV every 8 weeks.	_	
☐ Rinvoq tm	☐ 15 mg tablet	☐ Take 1 tablet by mouth o	nce daily with or without food.		
☐ Simponi®	□ 50 mg Autoinjector□ 50 mg Prefilled Syringe	☐ Inject 50 mg subQ once a month.			
☐ Simponi Aria®	□ 50 mg/4 mL Single-Use Vi Patient Dosing Wt: kg	maintenance dose.	□ Initial Dose: Infuse mg (2mg/kg) IV over 30 minutes at Weeks 0, 4, then maintenance dose. □ Maintenance Dose: Infuse mg (2mg/kg) IV over 30 minutes every 8 weeks.		
□ Stelara®	□ 45 mg/0.5 mL Prefilled Syr (PsA) □ 90 mg/mL Prefilled Syring (PsA with coexistent Psorias patients > 100 kg)	☐ <u>Initial Dose</u> : Inject one s e ☐ <u>Maintenance Dose</u> : Inje	☐ Initial Dose: Inject one syringe subQ at Weeks 0 and 4, then maintenance dose. ☐ Maintenance Dose: Inject one syringe subQ every 12 weeks.		
□ Taltz® □ 80 mg Autoinjector □ 80 mg Prefilled Syringe		maintenance dose startir	□ Initial Dose: Inject 160 mg (two 80 mg injections) subQ on Day 1, then maintenance dose starting Day 29. □ Maintenance Dose: Inject 80 mg subQ every 4 weeks.		
☐ Xeljanz [®] ☐ 5 mg tablet		☐ Take one tablet by mouth	☐ Take one tablet by mouth twice daily.		
☐ Xeljanz® XR ☐ 11 mg tablet		☐ Take one tablet by mouth	n once daily.		
Deliver te: Patient Office Other: Date: Needs by Date:					
Prescriber Authorization: I authorize this pharmacy and its representatives to act as my authorized agent to secure coverage and initiate the insurance prior authorization process for my patient(s), and to sign any necessary forms on my behalf as my authorized agent, including the receipt of any required prior authorization forms and the receipt and submission of patient lab values and other patient data. In the event that this pharmacy determines that it is unable to fulfill this prescription, I further authorize this pharmacy to forward this information and any related materials related to coverage of the product to another pharmacy of the patient's choice or in the patient's insurer's provider network. □ Product Substitution Permitted □ Dispense as Written					